

Subthreshold cascade production in heavy ion collisions

L. Feng, L.W. Chen,¹ C.M. Ko, and S.H. Lee²

¹INPAC, Department of Physics and Shanghai Key Laboratory for Particle Physics and Cosmology, Shanghai Jiao Tong University, Shanghai 200240, China

²Institute of Physics and Applied Physics, Yonsei University, Seoul 120-749, Korea

We have calculated the cross sections for the reaction $YY \rightarrow N\Xi$ ($Y = \Lambda, \Sigma$) based on a gauged SU(3)-invariant hadronic Lagrangian in the Born approximation [1] and found that these cross sections are almost four times the cross sections for the reaction $KY \rightarrow \pi\Xi$ that was considered in a previous study [2]. We then used these cross sections to study Ξ production in $^{40}\text{Ar}+^{40}\text{KCl}$ collisions at the subthreshold energy of 1.76 AGeV within the framework of a relativistic transport model that includes explicitly the nucleon, delta, pion, and perturbatively the kaon, antikaon, hyperons, and Ξ [3]. We found that the reaction $YY \rightarrow N\Xi$ would enhance the abundance by a factor of about 16 compared to that from the reaction $KY \rightarrow \pi\Xi$, resulting in an abundance ratio $\Xi^-/(\Lambda+\Sigma^0) = 3.38 \times 10^{-3}$ that is essentially consistent with that measured by the HADES Collaboration at GSI [4]. Our study has thus helped in resolving one of the puzzles in particle production from heavy ion collisions at subthreshold energies.




FIG. 1. Left window: Cross sections for (a) $\Lambda\Lambda \rightarrow N\Xi$, (b) $\Lambda\Sigma \rightarrow N\Xi$, (c) $\Sigma\Sigma \rightarrow N\Xi$, (d) $N\Xi \rightarrow \Lambda\Lambda$, (e) $N\Xi \rightarrow \Lambda\Sigma$, and (f) $N\Xi \rightarrow \Sigma\Sigma$ as functions of the center-of-mass energy from the Born approximation with cutoff parameters $\Lambda = 0.5$ GeV (dashed lines), $\Lambda = 0.7$ GeV (solid lines), and $\Lambda = 1$ GeV (dotted lines). Right window: Time evolutions of (a) central baryon density (right scale) and the abundances (left scales) of π and Δ , (b) K , Λ , Σ , and antikaon, and (c) Ξ produced from different reactions.

- [1] F. Li, L.W. Chen, C.M. Ko, and S. H. Lee, Phys. Rev. C **85**, 064902 (2012).
- [2] C.H. Li and C.M. Ko Nucl. Phys. **A712**, 110 (2002).
- [3] C.M. Ko and G.Q. Li, J. Phys. G **22**, 1673 (1996).
- [4] N. Kaiser and W. Weise, Phys. Lett. B **512**, 283 (2001).