

THE NATURE OF LIGHT DARK MATTER

Peter L. Biermann^{1,2,3}, Alexander Kusenko⁴,
Faustin Munyaneza¹ &
Jaroslaw Stasielak^{1,2,5,6}

¹ MPI for Radioastronomy, Bonn, Germany

² Dep. of Phys. & Astron., Univ. of Bonn

³ Dep. of Phys. & Astr., Univ. of Alabama,
Tuscaloosa, AL, USA

⁴ Dep. of Phys. & Astr., Univ. of Cal.,
Los Angeles, CA, USA

⁵ Physics Dept., Jagell. Univ., Cracow

⁶ KASI, Daejeon, Korea

www.mpifr-bonn.mpg.de/div/theory

Dark Matter: The evidence

- Dark Matter is required:
- for stability of disks in galaxies (e.g. Ostriker)
- to explain the rotation curves (e.g. Rubin)
- to explain the hot gas distribution in early Hubble type galaxies (e.g. Biermann)
- to explain the motions in groups and clusters (e.g. Zwicky)
- to explain the hot gas containment in groups and clusters (e.g. Ensslin et al.)
- to explain the structure formation
- to explain the flat geometry of the Universe (Spergel et al. 2003, 2007)

70 years: What is dark matter?

Dark Matter: Some history

Sterile neutrinos? Weakly interacting neutrinos - interact with other neutrinos and gravity

- 1930ies: "Dark Matter" first: Oort (1932), Zwicky (1933, 1937), and Swift (1936)
- 1970: Pontecorvo considers sterile neutrinos and introduces the name
- 1982: Olive & Turner consider sterile neutrinos in general terms
- 1994: Dodelson & Widrow show that sterile neutrinos can be dark matter
- 1997: Kusenko & Segre show that sterile neutrinos can explain pulsar kicks
- 2006: Shaposhnikov: The various origins of sterile neutrinos

Proposal

- Lightest supersymmetric particle, very massive: problem small scales
- However: G. Gilmore et al find many small dwarf galaxies, and smaller satellites are hindered in formation by ionization
- Sterile neutrinos at $2 \text{ keV} \lesssim m_{DM} \lesssim 20 \text{ keV}$.
- Right handed neutrinos, very weakly interact with normal left handed neutrinos; otherwise interact only via gravitation
- Lifetime $\tau = 1.3 \cdot 10^{26} \text{ s} \left(\frac{7 \text{ keV}}{m_{DM} c^2} \right)^5 \left(\frac{0.8 \cdot 10^{-9}}{\sin^2 \theta} \right)$.
- They decay, most importantly: one active neutrino and a photon
- Never in thermodynamic equilibrium, require a chemical potential; probably far sub-thermal

Background

- They can be produced in the right amount (Dodelson & Widrow 1994; Shi & Fuller 1999; Abazajian *et al.* 2001; Dolgov & Hansen 2002; Abazajian 2006)
- They can explain the baryon asymmetry (Akhmedov *et al.* 1998; Asaka & Shaposhnikov 2005)
- They can explain the lack of power on small scales in Large Scale Structure Formation (Olive & Turner 1982; Abazajian 2006)
- They can explain the inner dark matter distribution and spatial shape in galaxies (Belokurov *et al.* 2006; Fellhauer *et al.* 2006)

Lots of recent work by Kevork Abazajian and Mikhail Shaposhnikov

Our recent work

- Pulsar kicks, velocities up to 1000 km/s – see the guitar nebula – (Kusenko 2004)
range: 2 to 20 keV
- ! Transition from active neutrino to weakly interacting neutrino to prevent scattering: right handed neutrino!
- Early growth of black holes (Munyanenza & Biermann 2005)
range: 12 - 450 keV: ! Fermion
- Aspen September 2005: Overlap - Eureka!
- From increased secondary ionization molecular Hydrogen formed more abundantly, and so star formation triggered early (Biermann & Kusenko 2007; Stasielak, Biermann & Kusenko 2006) – this removed major obstacle (Yoshida *et al.* 2003) for this known proposal

First star formation

- Sterile neutrinos decay into an active neutrino and a photon (of half the mass in energy)
- After redshift about 40 no absorption anymore, before fraction of ionized atoms (η fraction of photon energy into ionization)
$$x_e^{(s)} \sim \frac{0.2}{(1+z)^{3/2}} \left(\frac{\eta}{0.3}\right) \left(\frac{m_s}{7 \text{ keV}}\right)^5 \left(\frac{\sin^2 \theta}{0.8 \cdot 10^{-9}}\right)$$
- Formation of H_2 via H^-
- Level of H_2 above $5 \cdot 10^{-4}$ (Tegmark et al. 1997) strong cooling
- Almost an order of magnitude enhancement, with maximum at redshift 80
- Corresponding heating cancels effect for redshifts < 20 (Mapelli, Ripamonti 2006)

The first stars

- Star formation triggered early
- Early magnetic fields from first massive stars – the fastest process – then cosmic ray driven dynamo process in galactic disks, implies by necessity galactic winds
- Reionization from first massive stars at $z > 10$
- Energy input from stellar winds and supernovae
- Chemical enrichment from first massive stars (Wolf Rayet star winds and supernovae)
- First dust
- First Gamma Ray Bursts
- First stellar black holes
- First cosmic rays

Primordial magnetic fields

- Ionization by secondary electron, from ionization:
- The time scale for energy gain (e.g., Jokipii 2004)

$$\tau_A = \frac{4\kappa}{V_A^2} \quad (1)$$

- Assuming $\frac{B^2/8\pi}{I(k)k} \simeq 1$ and $I(k) \sim k^{-1}$ to just cancel the energy loss by ionization $B(z = 100) < 2 \cdot 10^{-8}$ Gauß, corresponding to today

$$B < 4 \cdot 10^{-12} \text{ Gauß.} \quad (2)$$

- K. Dolag et al. (2002, 2005):
 $B < (0.2 - 1.) \times 10^{-12}$ Gauß from structure formation simulations: consistent !

Galaxies

- Galaxies merge, and then
- Energy transported out, matter transported in, due to extreme mixing in a merger; akin to accretion disk theory (Lüst 1952)
- Inner dark matter distribution: Powerlaw, Moore 1998, Klypin et al. 2002, Navarro, Frenk, & White 1997
- Consider the low momentum tail of phase space distribution; for keV Fermions degenerate configuration: DARK MATTER STAR
- Suggestion, that Galactic Center Black Hole possibly only fed from dark accretion
- Alternative: direct collapse from supermassive star, or very compact stellar cluster: about $10^6 M_{\odot}$

X-ray background

- 2005 - 2007: Boyarsky et al., Riemer-Sørensen, Watson, ...
- What is the upper limit to an invisible emission line at an energy $m_{DM}c^2/2$
- X-ray background: if dark matter, then $m_{DM} < 10$ keV
- Neighboring clusters of galaxies (Virgo, Coma)
- Neighboring galaxies (M31)
- With major effort may be positively detectable with the X-ray satellites Chandra and Newton: Large field high spectral resolution spectroscopy
- Neighboring dwarf ellipticals: Suzaku observations: M. Loewenstein, Kusenko, PLB

Lyman- α forest

- McDonald et al. 2001: forest example; simulations, weak fluctuations in density
- Seljak et al. 2006: WDM power spectrum versus CDM power spectrum
- McDonald et al. 2000: Contamination by metal lines?
- Seljak et al. 2006: SDSS data and high resolution data versus models in CDM (thick lines) and WDM (thin lines)
- Seljak et al. 2006: Discrepancy at high redshift: $m_{DM} > 14$ keV
- Viel et al. 2006: same analysis $m_{DM} > 10$ keV
- Reconciliation with lower mass by subthermal property of particle?
→ $m_{DM} > 2$ keV

Galaxies

Work of group around G. Gilmore

- Shape of our Galaxy halo: Very spherical: From precessional disruption of incoming dwarf galaxies
- Mass of dwarf spheroidal galaxies
- All data consistent with lower limit to dark matter mass of $5 \cdot 10^7 M_{\odot}$, for a large range of baryonic mass - direct indication for warm dark matter (or formation history? effect of ionization, heating and cooling of satellite galaxy?)
- Based on Gilmore et al. (2007), and Wyse & Gilmore (2007):

$$m_{DM} < 4 \text{ keV (AK)}$$

Summary

- Right-handed neutrinos? far subthermal?
- First star formation; first reionization; first magnetic fields; first cosmic rays; first black holes; first ultra high energy cosmic rays
- X-ray line from galaxies, and background
- Pulsar kicks
- Lyman α -forest - consistent for subthermal distribution
- Galactic halo structure; dwarf spheroidal galaxies
- All of these findings can be explained independently. If one concept: sterile neutrino
- $\sim 2 \text{ keV} \lesssim m_s \lesssim \sim 4 \text{ keV}$
- **Critical test X-ray emission line**
Weakly Interacting Neutrinos

Acknowledgements

Support for PLB is coming from the AUGER membership and theory grant 05 CU 5PD 1/2 via DESY/BMBF. Support for AK is coming from DOE grant DE-FG03-91ER40662 and NASA ATP grants NAG 5-10842 and NAG 5-13399. Support for FM is coming from the Humboldt Foundation.

References

- [1] “Observed properties of dark matter on small spatial scales”, Wyse, & Gilmore (2007); arXiv/0708.1492
- [2] “The observed properties of dark matter on small scales”, Gilmore et al. (2007); astro-ph/0703308

- [3] “Constraints on Sterile Neutrino Dark Matter”, Abazajian, K., Koushiappas, S. M., *Phys. Rev. D* (submitted, 2006), astro-ph/0605271
- [4] “Linear cosmological structure limits on warm dark matter”, Abazajian, K., *Phys. Rev. D* **73**, id. 063513 (2006), astro-ph/0512631
- [5] “Production and evolution of perturbations of sterile neutrino dark matter”, Abazajian, K., *Phys. Rev. D* **73**, id. 063506 (2006), astro-ph/0511630
- [6] “Cosmological lepton asymmetry, primordial nucleosynthesis and sterile neutrinos”, Abazajian, K., *Phys. Rev. D* **72**, id. 063004 (2005)
- [7] “Sterile neutrino hot, warm, and cold dark matter”, Abazajian, K., Fuller, G.

- M., Patel, M., *Phys. Rev. D* **64**, id. 023501 (2001)
- [8] “Baryogenesis via Neutrino Oscillations”, Akhmedov, E. Kh., Rubakov, V. A., Smirnov, A. Yu., *Phys. Rev. Letters* **81**, 1359-1362 (1998)
- [9] “The ν MSM, dark matter and baryon asymmetry of the universe, Asaka, T., Shaposhnikov, M., *Phys. of Fluids* **620**, 17-26 (2005)
- [10] “Opening a new window for warm dark matter”, Asaka, T., Shaposhnikov, M., Kusenko, A., *Phys. of Fluids* **638**, 401-406 (2006), hep-ph/ 0602150
- [11] “The ν MSM, dark matter and neutrino masses”, Asaka, T., Blanchet, St., Shaposhnikov, M., *Phys. of Fluids* **631**, 151-156 (2005)

- [12] “The Field of Streams: Sagittarius and Its Siblings”, Belokurov, V. *et al.*, *Astrophys. J. Letters* **642**, L137 - L140 (2006), astro-ph/0605025)
- [13] “The detection of hot intergalactic gas in the NGC 3607 group of galaxies with the Einstein satellite”, Biermann, P., Kronberg, P. P.; Madore, B. F., *Astrophys. J. Letters* **256**, p. L37-L40 (1982)
- [14] “Detection of 10 to the 10th solar masses of hot gas in the normal elliptical galaxy NGC 5846 with the Einstein satellite”, Biermann, P., Kronberg, P. P., *Astrophys. J. Letters* **268**, p. L69-L73 (1983)
- [15] “Relic keV sterile neutrinos and reionization”, P.L. Biermann & A. Kusenko, 2006, *Phys. Rev. Letters* **96**, 091301 (2006); astro-ph/0601004

- [16] “Ultra high energy cosmic rays from sequestered X bursts [rapid communication]”, P.L. Biermann & P. Frampton 2006, *Physics Letters B*, **634**, p. 125-129 (2006); astro-h/0512188
- [17] “Cosmic-ray protons and magnetic fields in clusters of galaxies and their cosmological consequences”, Torsten A. Enßlin, Peter L. Biermann, Philipp P. Kronberg, and Xiang P. Wu, *Astrophys. J.* **477**, 560 (1997); astro-ph/9609190
- [18] “Where to find a dark matter sterile neutrino?”, Boyarsky, A., Neronov, A., Ruchayskiy, O., Shaposhnikov, M., Tkachev, I., astro-ph/0603660
- [19] “Restrictions on parameters of sterile neutrino dark matter from observations of galaxy clusters”, Boyarsky, A.,

- Neronov, A., Ruchayskiy, O., Shaposhnikov, M., astro-ph/0603368
- [20] “Constraints on sterile neutrino as dark matter candidate from the diffuse X-ray background”, A. Boyarsky, A. Neronov, O. Ruchayskiy, M. Shaposhnikov, *Month. Not. Roy. Astr. Soc.* in press, July 2006, astro-ph/0512509
- [21] “Sterile neutrinos as dark matter”, Dodelson, S., Widrow, L. M., *Phys. Rev. Letters* **72**, 17-20 (1994)
- [22] “Massive sterile neutrinos as warm dark matter”, Dolgov, A. D., Hansen, S. H., *Astropart. Phys.* **16**, 339-344 (2002)
- [23] “The Origin of the Bifurcation in the Sagittarius Stream”, Fellhauer, M., *et al.*, (2006), astro-ph/0605026
- [24] Kusenko & Segre (1997)

- [25] “Pulsar kicks from neutrino oscillations,” A. Kusenko, *Int. J. Mod. Phys. D* **13**, 2065 (2004); astro-ph/0409521
- [26] “Cosmological lower bound on heavy-neutrino masses”, Lee, B. W., Weinberg, St., *Phys. Rev. Letters* **39**, 1977, pp.165-168
- [27] “Fast Growth of supermassive black holes in Galaxies”, F. Munyaneza & P.L. Biermann, *Astron. & Astroph.* **436**, 805 - 815 (2005); astro-ph/0403511
- [28] “Cosmological bounds on the masses of stable, right-handed neutrinos”, Olive, K. A., Turner, M. S., *Phys. Rev.* **D 25**, 213 - 216 (1982)
- [29] Oort, J.H., *Bull. Astron. Inst. Netherl.*, **VI**, 249 (1932)
- [30] Pontecorvo (1970)

- [31] “The impact of dark matter decays and annihilations on the formation of the first structures”, E. Ripamonto, M. Mapelli, A. Ferrara, *Month. Not. Roy. Astr. Soc.* (2006), astro-ph/...
- [32] “Can sterile neutrinos be the dark matter?”, U. Seljak, A. Makarov, P. McDonald, H. Trac, *Phys. Rev. D* (2006), astro-ph/0602430
- [33] “Cosmological parameters from combining the Lyman- α forest with CMB, galaxy clustering and SN constraints”, U. Seljak, A. Slosar, P. McDonald, , *Phys. Rev. D* (2006), astro-ph/0604335
- [34] “The ν MSM, dark matter and neutrino masses”, Shaposhnikov, M., *Journal of Physics: Conference Series*, Volume 39, Issue 1, pp. 176-178 (2006).

- [35] “The ν MSM, dark matter and baryon asymmetry of the Universe”, Shaposhnikov, M., *Journal of Physics: Conference Series*, **39**, 9-11 (2006).
- [36] “The ν MSM, Inflation, and Dark Matter”, Shaposhnikov, M., Tkachev, I., (2006), hep-ph/0604236
- [37] “New Dark Matter Candidate: Nonthermal Sterile Neutrinos”, Shi, X. D., Fuller, G. M., *Phys. Rev. Letters* **82**, 2832 - 2835 (1999)
- [38] “First-Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Determination of Cosmological Parameters”, Spergel, D.N., *et al.*, *Astrophys. J. Suppl.* **148**, 175 (2003), astro-ph/0302209

- [39] “Wilkinson Microwave Anisotropy Probe (WMAP) Three Year Results: Implications for Cosmology”, Spergel, D.N., *et al.*, *Astrophys. J.* (submitted, 2006), astro-ph/0603449
- [40] “A Large Dark Matter Core in the Fornax Dwarf Spheroidal Galaxy?”, Strigari, L. E. *et al.*, (2006), astro-ph/0603775
- [41] “Can sterile neutrinos be ruled out as warm dark matter candidates?”, M. Viel, J. Lesgourgues, M. G. Haehnelt, S. Mattares, A. Riotto, *Phys. Rev. D* (2006), astro-ph/0605706
- [42] “Signatures for a Cosmic Flux of Magnetic Monopoles”, Stuart D. Wick, Thomas W. Kephart, Thomas J. Weiler, Peter L. Biermann, *Astro. Part. Phys.* **18**, 663 - 687 (2003), astro-ph/0001233

- [43] “Early Structure Formation and Reionization in a Warm Dark Matter Cosmology”, Yoshida, N., Sokasian, A., Hernquist, L., Springel, V., *Astrophys. J. Letters* **591**, L1 - L4. (2003)
- [44] Zwicky, F. *Helv. Phys. Acta* **6**, 110 (1933)
- [45] “On the masses of nebulae and of clusters of nebulae”, Zwicky, F., *Astrophys. J.* **86**, 217 (1937)