

# PHYS 606 - Spring 2014 - Homework X Solution

$$[1] a) [L_i, r_j] = \epsilon_{ike} [r_k p_e, r_j] = \epsilon_{ike} \left( r_k \underbrace{[p_e, r_j]}_{-i\hbar \delta_{je}} + \underbrace{[r_k, r_j]}_{=0} p_e \right) = i\hbar \epsilon_{ijk} r_k$$

$$[L_i, p_j] = \epsilon_{ike} \left( r_k \underbrace{[p_e, p_j]}_{=0} + \underbrace{[r_k, p_j]}_{i\hbar \delta_{kj}} p_e \right) = i\hbar \epsilon_{ije} p_e$$

$$[L_i, K_j] = [L_i, m r_j - p_j t] = i\hbar \epsilon_{ijk} (m r_k - p_k t) = i\hbar \epsilon_{ijk} K_k$$

$$[1] b) [L_{\pm}, L_z] = \underbrace{[L_x, L_z]}_{-i\hbar L_y} \pm i \underbrace{[L_y, L_z]}_{i\hbar L_x} = \mp i\hbar (L_x \pm iL_y) = \mp i\hbar L_{\pm}$$

$$[L_+, L_-] = i[L_y, L_x] - i[L_x, L_y] = 2i(-i\hbar L_z) = 2\hbar L_z$$

$$L^2 - L_z^2 = L_x^2 + L_y^2 = (L_x \pm iL_y)(L_x \mp iL_y) - L_x(\mp iL_y) - (\pm iL_y)L_x$$

$$= L_{\pm} L_{\mp} \pm i \underbrace{[L_x, L_y]}_{i\hbar L_z} = L_{\pm} L_{\mp} \mp i\hbar L_z$$

$$[2] a) \langle n' | a | n \rangle = C_n \langle n' | n-1 \rangle = C_n \delta_{n', n-1} \quad \forall n, n' \in \mathbb{N}$$

$$\langle n' | a^\dagger | n \rangle = D_n \delta_{n', n+1}$$

I.e. in explicit matrix form

$$a = \begin{pmatrix} 0 & c_1 & 0 & 0 & \dots \\ 0 & 0 & c_2 & 0 & \dots \\ 0 & 0 & 0 & c_3 & \dots \\ \dots & \dots & \dots & \dots & \dots \end{pmatrix} \quad a^\dagger = \begin{pmatrix} 0 & 0 & 0 & \dots \\ D_0 & 0 & 0 & \dots \\ 0 & D_1 & 0 & \dots \\ 0 & 0 & D_2 & \dots \\ \dots & \dots & \dots & \dots \end{pmatrix}$$

$$[2] b) |C_n|^2 = |\langle n-1 | a | n \rangle|^2 = \langle n | a^\dagger | n-1 \rangle \langle n-1 | a | n \rangle = \sum_{n' \in \mathbb{N}} \langle n | a^\dagger | n' \rangle \langle n' | a | n \rangle$$

$$= \langle n | a^\dagger a | n \rangle = n$$

all matrix elements for  $n' \neq n-1$  vanish

$$\Rightarrow C_n = \sqrt{n} \quad (\text{choose phase to be zero})$$

2

$$\text{Similarly } |D_n|^2 = |\langle n+1 | a^\dagger | n \rangle|^2 = \sum_{n' \in \mathbb{N}} \langle n | a | n' \rangle \langle n' | a^\dagger | n \rangle$$

$$= \langle n | a a^\dagger | n \rangle = \langle n | a^\dagger a + \mathbb{1} | n \rangle = n+1$$

$$\Rightarrow D_n = \sqrt{n+1}$$

$$|n\rangle = N_n (a^\dagger)^n |0\rangle = N_n \sqrt{n} \cdot \sqrt{n-1} \cdot \dots |0\rangle = N_n \sqrt{n!} |0\rangle$$

$$|n\rangle \text{ and } |0\rangle \text{ normalized to unity} \Rightarrow N_n = \frac{1}{\sqrt{n!}}$$

$$(c) \hat{x} = \sqrt{\frac{\hbar}{2m\omega}} (a + a^\dagger) \Rightarrow \hat{x} = \sqrt{\frac{\hbar}{2m\omega}} \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & \sqrt{2} & 0 \\ 0 & \sqrt{2} & 0 & \sqrt{2} \\ 0 & 0 & \sqrt{3} & 0 \\ \vdots & \vdots & \vdots & \vdots \end{pmatrix}$$

$$(d) \hat{x} |x\rangle = x |x\rangle \Rightarrow \langle n | \hat{x} |x\rangle = x \langle n | x\rangle \Rightarrow \sum_{n' \in \mathbb{N}} \langle n | \hat{x} | n' \rangle \langle n' | x\rangle = x \langle n | x\rangle$$

$$\Rightarrow \sum_{n' \in \mathbb{N}} \langle n | x | n' \rangle \psi_{n'}(x) = x \psi_n(x)$$

$$\text{from (c): } \langle n | x | n' \rangle = \left( \sqrt{n+1} \delta_{n, n'-1} + \sqrt{n} \delta_{n, n'+1} \right) \sqrt{\frac{\hbar}{2m\omega}}$$

$$\Rightarrow \sqrt{n+1} \psi_{n+1} + \sqrt{n} \psi_{n-1} = x \psi_n(x) \sqrt{\frac{2m\omega}{\hbar}} \quad (\text{for } n=0: \psi_{-1} = 0) \quad (\Delta)$$

$$(e) \text{ Ansatz } \psi_n(x) = 2^{-\frac{n}{2}} (n!)^{-\frac{1}{2}} h_n \left( \sqrt{\frac{m\omega}{\hbar}} x \right) \psi_0(x)$$

$$\text{with } h_0 = 1 \text{ and } h_{n+1}(x) - 2x h_n(x) + 2n h_{n-1}(x) = 0 \quad (\square)$$

$$\text{into } (\Delta): \sqrt{n+1} 2^{-\frac{n+1}{2}} (n+1)!^{-\frac{1}{2}} h_{n+1} \left( \sqrt{\frac{m\omega}{\hbar}} x \right) \psi_0(x) + \sqrt{n} 2^{-\frac{n-1}{2}} (n-1)!^{-\frac{1}{2}} h_{n-1} \left( \sqrt{\frac{m\omega}{\hbar}} x \right) \psi_0(x)$$

$$= x \sqrt{\frac{2m\omega}{\hbar}} 2^{-\frac{n}{2}} (n!)^{-\frac{1}{2}} h_n \left( \sqrt{\frac{m\omega}{\hbar}} x \right) \psi_0(x)$$

$$\text{After dividing by common factors: } \frac{1}{2} \sqrt{\frac{\hbar}{m\omega}} \frac{1}{\sqrt{(n+1)n}} h_{n+1}(\xi) + \sqrt{n} h_{n-1}(\xi) = \xi \frac{\sqrt{2}}{\sqrt{2}\sqrt{n}} h_n(\xi)$$

$$\text{define } \xi = \sqrt{\frac{m\omega}{\hbar}} x$$

$$\Rightarrow h_{n+1}(\xi) - 2\xi h_n(\xi) + 2n h_{n-1}(\xi) = 0$$

as in  $(\square)$ .

(f) Use  $H_n(\xi) = (-1)^n e^{\xi^2} \frac{d^n}{d\xi^n} e^{-\xi^2}$  (HW III, [1])

$$\begin{aligned} H_{n+1}(\xi) &= (-1)^{n+1} e^{\xi^2} \frac{d^{n+1}}{d\xi^{n+1}} e^{-\xi^2} = (-1)^{n+1} e^{\xi^2} \left( -2n \frac{d^n}{d\xi^n} - 2\xi \frac{d^n}{d\xi^n} \right) e^{-\xi^2} \\ &= -2n H_n(\xi) + 2\xi H_n(\xi) \quad \text{which is the relation from (3).} \end{aligned}$$

[3] (a) From V.4 in the lecture notes we already know that the radial

equation for  $R(r)$  in the case  $V(\vec{r}) = 0$  is

$$\left[ -\frac{\hbar^2}{2m} \frac{1}{r^2} \frac{d}{dr} \left( r^2 \frac{d}{dr} \right) + \frac{l(l+1)\hbar^2}{2mr^2} \right] R(r) = ER(r)$$

Introduce  $\rho = \frac{r}{\hbar} \sqrt{2mE} = kr$  with  $k = \frac{1}{\hbar} \sqrt{2mE}$  (magnitude of the wave vector of the free particle)

$$\frac{1}{\rho^2} \frac{d}{d\rho} \left( \rho^2 \frac{d}{d\rho} \right) R(\rho) - \frac{l(l+1)}{\rho^2} R(\rho) + R(\rho) = 0$$

$$\Rightarrow \frac{d^2}{d\rho^2} R + \frac{2}{\rho} \frac{dR}{d\rho} + \left( 1 - \frac{l(l+1)}{\rho^2} \right) R = 0 \quad (*)$$

(b) Ansatz  $j_l(\rho) = \frac{\rho^e}{2^{e+1} e!} \int_{-1}^{+1} e^{i\rho s} (1-s^2)^e ds$  (also known as Poisson's integral representation for sph. Bessel fcts.)

into (\*): [dropping common factors]

$$\begin{aligned} &\rho^e \int_{-1}^{+1} (i\rho s)^2 e^{i\rho s} (1-s^2)^e ds + 2\rho^e \int_{-1}^{+1} (i\rho s) e^{i\rho s} (1-s^2)^e ds + l(l-1) \rho^{e-2} \int_{-1}^{+1} e^{i\rho s} (1-s^2)^e ds \\ &+ 2\rho^e \int_{-1}^{+1} (i\rho s) e^{i\rho s} (1-s^2)^e ds + 2l \rho^{e-2} \int_{-1}^{+1} e^{i\rho s} (1-s^2)^e ds + \rho^e \int_{-1}^{+1} e^{i\rho s} (1-s^2)^e ds \\ &- l(l+1) \rho^{e-2} \int_{-1}^{+1} e^{i\rho s} (1-s^2)^e ds = 0 \end{aligned}$$

$$\Rightarrow \rho^e \int_{-1}^{+1} e^{i\rho s} (1-s^2)^{e+1} ds + 2(l+1) \rho^{e-1} \int_{-1}^{+1} (i\rho s) e^{i\rho s} (1-s^2)^e ds = 0$$

$$= -2\rho^{e-1} \int_{-1}^{+1} e^{i\rho s} \frac{d}{ds} (1-s^2)^{e+1} ds$$

$$\Rightarrow 0 = 0$$

i.e. the  $j_l$  are solutions.

$$= \underbrace{e^{i\rho s} (1-s^2)^{e+1}}_{\text{part. ind.}} \Big|_{-1}^{+1} - \int_{-1}^{+1} i\rho s e^{i\rho s} (1-s^2)^{e+1} ds = 0$$

4

[c] Induction! Check  $l=0$ :  $\frac{\sin s}{s} = \frac{1}{2} \int_{-1}^{+1} \cos(s\xi) d\xi = \frac{1}{2} \int_{-1}^{+1} e^{is\xi} d\xi = j_0 \quad \checkmark$

Suppose we know  $(-1)^l s^l e^{\left(\frac{d}{ds}\right)^l \frac{\sin s}{s}} = j_l(s)$ ; then for  $l+1$ :

$$j_{l+1}(s) = \frac{s^{l+1}}{2^{l+2}(l+1)!} \int_{-1}^{+1} e^{is\xi} (1-\xi^2)^{l+1} d\xi = \frac{s^{l+1}}{2^{l+2}(l+1)!} \int_{-1}^{+1} (l+1) \frac{-2\xi}{i s} e^{is\xi} (1-\xi^2)^l d\xi$$

$$= \frac{s^l}{2^{l+1} l!} \int_{-1}^{+1} \left(-\frac{d}{ds} e^{is\xi}\right) (1-\xi^2)^l d\xi \quad + \text{boundary term } \alpha(1-\xi^2)^l \Big|_{\xi=\pm 1} = 0$$

$$= (-1)^{l+1} s^l e^{\frac{d}{ds} \left(\frac{d}{ds}\right)^l \frac{\sin s}{s}} = (-1)^{l+1} s^{l+1} \left(\frac{d}{ds}\right)^{l+1} \frac{\sin s}{s}$$

Explicitly:  $j_0(s) = \frac{\sin s}{s} \quad j_1(s) = \frac{\sin s}{s^2} - \frac{\cos s}{s}$

$$j_2(s) = \frac{3\sin s}{s^3} - \frac{3\cos s}{s^2} - \frac{\sin s}{s}$$

Plots: see attached

[d] Plane wave in  $z$ -direction in spherical coordinates:

$$e^{i\vec{k}\cdot\vec{r}} = e^{ikr\cos\theta} = \sum_{l=0}^{\infty} \sum_{m=-l}^{+l} C_{lm} j_l(kr) Y_l^m(\theta, \phi) \quad (\text{completeness of states in sph. coord.})$$

$$= \sum_{l=0}^{\infty} C_l j_l(kr) P_l(\cos\theta) \quad (\text{since } \phi \text{ does not occur on l.h.s. } m=0!)$$


$$\Rightarrow \int_{\xi=\cos\theta}^{+1} e^{ikr\xi} P_l(\xi) d\xi = \sum_{l=0}^{\infty} C_l j_l(kr) \int_{-1}^{+1} P_l(\xi) P_l(\xi) d\xi = \frac{2}{2l+1} C_l j_l(kr)$$

$$\Rightarrow C_l = (2l+1) \frac{1}{2j_l(kr)} \int_{-1}^{+1} e^{ikr\xi} P_l(\xi) d\xi = \frac{(2l+1)}{j_l(kr)} \frac{1}{2^{l+1} l!} \int_{-1}^{+1} \underbrace{\left(\frac{d}{d\xi} e^{ikr\xi}\right)}_{\substack{\text{part. int.} \\ l \text{ times}}} (1-\xi^2)^l d\xi$$


$$= (2l+1) i^l \quad + \text{boundary terms which vanish} = i^l j_l(kr)$$

$$\Rightarrow e^{ikr\cos\theta} = \sum_{l=0}^{\infty} (2l+1) i^l j_l(kr) P_l(\cos\theta)$$

{SphericalBesselJ[n,x], n =0}



{SphericalBesselJ[n,x], n =1}



{SphericalBesselJ[n,x], n =2}

