

---

---

# Physics 606 — Spring 2015

## Homework 2

Instructor: Rainer J. Fries

Turn in your work by February 17

---

---

### [1] Gaussian Wave Packets — Part II (25 points)

At time  $t = 0$  consider a Gaussian wave packet (cf. HW I, [3]) centered around  $x_0$  with an average momentum  $k_0$ , i.e.

$$\psi(x, 0) = \left(\sqrt{2\pi}\sigma\right)^{-1/2} e^{ik_0x} e^{-\frac{(x-x_0)^2}{4\sigma^2}} \quad (1)$$

- If this packet represents a free particle of energy  $E = k_0^2/2m$  calculate the time evolution  $\psi(x, t)$  in explicit form.
- Using a program of your choice (Mathematica, Matlab, etc.) plot the real and imaginary parts of  $\psi(x, t)$  as well as  $|\psi(x, t)|^2$  as functions of  $x$  for different values of  $t$ . Choose suitable parameters to document the spreading of the wave packet with time. What determines the “speed” with which the width of the wave packet increases?

### [2] Schrödinger Equation in Momentum Space (20 points)

Starting from the Schrödinger equation in coordinate space for a particle of mass  $m$  and potential energy  $V(\vec{r})$  explicitly show that

$$i\hbar \frac{\partial}{\partial t} \phi(\vec{p}, t) = \frac{p^2}{2m} \phi(\vec{p}, t) + V(i\hbar \nabla_p) \phi(\vec{p}, t) \quad (2)$$

where  $\phi(\vec{p}, t)$  is the Fourier transformation of a coordinate space solution  $\psi(\vec{r}, t)$ .

### [3] Continuity Equation (30 points)

- Consider a system of classical particles described by a density  $\rho(\vec{r}, t)$  and a velocity field  $\vec{v}(\vec{r}, t)$ . Consider a volume  $V \equiv V(t)$  co-moving with the particle flow, i.e. the particle number in this volume is conserved,

$$\frac{d}{dt} \int_{V(t)} \rho d^3r = 0. \quad (3)$$

From this condition derive the continuity equation

$$\frac{\partial}{\partial t} \rho + \nabla \cdot (\rho \vec{v}) = 0. \quad (4)$$

- Consider wave functions  $\psi_1(\vec{r}, t)$  and  $\psi_2(\vec{r}, t)$  which obey the same time-dependent Schrödinger equation. Derive the current density  $\vec{j}_{12}(\vec{r}, t)$  which fulfills the continuity equation

$$\frac{\partial}{\partial t} (\psi_1 \psi_2^*) + \nabla \cdot \vec{j}_{12} = 0. \quad (5)$$

- (c) Hence what is the current  $\vec{j}$  that satisfies the continuity equation for the probability density  $\rho = |\psi|^2$  of a single field  $\psi$ ? Show that  $\vec{j}$  goes towards the result from (a), i.e.  $\vec{j} \rightarrow \rho\vec{v}$  in the classical limit.

[4] **Discontinuous Potential Energy** (25 points)

Consider a potential energy function  $V(\vec{r})$  which is discontinuous at the plane  $z = 0$ . Show that wave functions with the boundary conditions

$$\lim_{\epsilon \rightarrow 0} \psi(x, y, -\epsilon, t) = \lim_{\epsilon \rightarrow 0} \psi(x, y, +\epsilon, t) \quad (6)$$

$$\lim_{\epsilon \rightarrow 0} \frac{\partial}{\partial z} \psi(x, y, -\epsilon, t) = \lim_{\epsilon \rightarrow 0} \frac{\partial}{\partial z} \psi(x, y, +\epsilon, t) \quad (7)$$

for each point  $(x, y, 0)$  on the surface and for all times  $t$  satisfy the continuity equation of quantum mechanics. In other words, both the wave function and its derivative normal to the surface have to be continuous.