

---

---

# Physics 606 — Spring 2015

## Homework 9

Instructor: Rainer J. Fries

Turn in your work by April 21

---

---

[1] **Legendre Polynomials and Legendre Functions** (40 points)

Consider the differential equation

$$\frac{d}{d\xi} \left( (1 - \xi^2) \frac{dP}{d\xi} \right) - \frac{m^2}{1 - \xi^2} P + \lambda P = 0 \quad (1)$$

for a function  $P(\xi)$ ,  $-1 < \xi < 1$ , with parameters  $m \in \mathbb{N}$  and  $\lambda \in \mathbb{R}$ . It is called *Legendre's differential equation*.

- (a) Consider the special case  $m = 0$ . Make a power series ansatz for the solution,  $P(\xi) = \sum_{j=1}^{\infty} a_j \xi^j$ . From the differential equation derive a recursion relation between coefficients  $a_j$  and  $a_{j+2}$ . Show that the power series diverges at the endpoints  $\xi = \pm 1$  **unless**  $\lambda = l(l + 1)$  where  $l \in \mathbb{N}$  is a non-negative integer.
- (b) The outcome of (a) suggests that the only physically acceptable, non-singular solutions to Legendre's equation for  $m = 0$  are polynomials and they can be labeled by a quantum number  $l$  with  $\lambda = l(l + 1)$ . Show that these *Legendre polynomials* are given by

$$P_l(\xi) = \frac{1}{2^l l!} \frac{d^l}{d\xi^l} (\xi^2 - 1)^l. \quad (2)$$

(The normalization is simply a convention.) What is the degree of  $P_l$ ?

- (c) Write down the first four Legendre polynomials ( $l = 0, 1, 2, 3$ ) explicitly.
- (d) Show that Legendre polynomials are mutually orthogonal with respect to a scalar product defined as integration over the interval  $[-1, 1]$ , and their norm is  $\sqrt{2/(2l + 1)}$ , i.e.

$$\int_{-1}^1 P_l(\xi) P_{l'}(\xi) d\xi = \frac{2}{2l + 1} \delta_{ll'}. \quad (3)$$

- (e) Now we return to the general case of Legendre's differential equation. Show that for  $m \leq l$  the functions

$$P_l^m(\xi) = (1 - \xi^2)^{\frac{m}{2}} \frac{d^m}{d\xi^m} P_l(\xi) \quad (4)$$

are solutions to (1). They are called *associated Legendre functions of the first kind*.

[2] **Angular Momentum Operators** (40 points)

- (a) Show the following commutation relations for the angular momentum operator  $\vec{L} = \vec{r} \times \vec{p}$ : (i)  $[L_j, L_k] = \epsilon_{jkl} i\hbar L_l$ ,  $j, k, l = 1, 2, 3$  where  $\epsilon_{jkl}$  is the usual anti-symmetric Levi-Civita tensor with  $\epsilon_{123} = 1$ ; (ii)  $[L_j, L^2] = 0$  for  $j = 1, 2, 3$  where  $L^2 = L_1^2 + L_2^2 + L_3^2$ .
- (b) Derive the nabla operator  $\nabla$  and the Laplace operator  $\Delta$  in spherical coordinates  $r, \theta, \phi$ .
- (c) Give explicit expressions of the operators  $L_x, L_y$  and  $L_z$ , in coordinate space representations in *spherical coordinates* and show that in particular

$$L^2 = -\hbar^2 \left[ \frac{1}{\sin^2 \theta} \frac{\partial^2}{\partial \phi^2} + \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left( \sin \theta \frac{\partial}{\partial \theta} \right) \right]. \quad (5)$$

- (d) Since  $L_z$  and  $L^2$  are commuting operators we can find common eigenfunctions. Solve the two eigenvalue equations<sup>1</sup>

$$L_z Y(\theta, \phi) = m\hbar Y(\theta, \phi), \quad (6)$$

$$L^2 Y(\theta, \phi) = \lambda\hbar^2 Y(\theta, \phi) \quad (7)$$

by choosing a separation ansatz  $Y(\theta, \phi) = \Phi(\phi)\Theta(\theta)$ . The functions  $Y(\theta, \phi)$  in proper normalization (discussed later) are called *spherical harmonics*.

*Hint: First solve for  $\Phi(\phi)$  (what are the allowed values for  $m$ ?) and then show that the equation for  $\Theta$  reduces to Legendre's differential equation from problem [1].*

[3] **Ground State Splitting for the Double Harmonic Oscillator** (20 points)

Consider a particle of mass  $m$  in a double harmonic oscillator potential  $V(x) = \frac{1}{2}m\omega^2(|x| - a)^2$  where  $a$  is the parameter determining the separation of the two harmonic oscillator minima.

- (a) We choose trial functions

$$\psi_{\pm}^n = N_{\pm}^n [\psi_n(x - a) \pm \psi_n(x + a)] \quad (8)$$

as discussed in section IV.4, where the  $\psi_n$  are the usual harmonic oscillator eigenfunctions. Calculate the values of the functional  $\langle H \rangle[\psi_{\pm}^0]$  for the case  $n = 0$ . As you know they are approximations to the energies of the true ground state and first excited state.

- (b) In the asymptotic limit  $a \rightarrow \infty$  the integrals you obtained in (a) should evaluate to simple expressions. Show that the leading terms in this limit are

$$\langle H \rangle[\psi_{\pm}^0] = \frac{1}{2}\hbar\omega \mp \frac{\alpha}{\sqrt{\pi}} e^{-\alpha^2} \quad (9)$$

in terms of the dimensionless parameter  $\alpha = \sqrt{m\omega/\hbar}x$ . Thus which trial function,  $\psi_+^0$  or  $\psi_-^0$ , is the approximation for the ground state?

---

<sup>1</sup>It is customary to write powers of  $\hbar$  explicitly in this eigenvalue problem.