

---

---

# Physics 606 — Spring 2017

## Homework 9

Instructor: Rainer J. Fries

Turn in your work by April 20

---

---

### [1] Legendre Polynomials and Legendre Functions (40 points)

Consider the differential equation

$$\frac{d}{d\xi} \left( (1 - \xi^2) \frac{dP}{d\xi} \right) - \frac{m^2}{1 - \xi^2} P + \lambda P = 0 \quad (1)$$

for a function  $P(\xi)$ ,  $-1 < \xi < 1$ , with parameters  $m \in \mathbb{N}$  and  $\lambda \in \mathbb{R}$ . It is called *Legendre's differential equation*.

- (a) Consider the special case  $m = 0$ . Make a power series ansatz for the solution,  $P(\xi) = \sum_{j=1}^{\infty} a_j \xi^j$ . From the differential equation derive a recursion relation between coefficients  $a_j$  and  $a_{j+2}$ . Show that the power series diverges at the endpoints  $\xi = \pm 1$  **unless**  $\lambda = l(l + 1)$  where  $l \in \mathbb{N}$  is a non-negative integer.
- (b) The outcome of (a) suggests that the only physically acceptable, non-singular solutions to Legendre's equation for  $m = 0$  are polynomials and they can be labeled by a quantum number  $l$  with  $\lambda = l(l + 1)$ . Show that these *Legendre polynomials* are given by

$$P_l(\xi) = \frac{1}{2^l l!} \frac{d^l}{d\xi^l} (\xi^2 - 1)^l. \quad (2)$$

(The normalization is simply a convention.) What is the degree of  $P_l$ ?

- (c) Write down the first four Legendre polynomials ( $l = 0, 1, 2, 3$ ) explicitly.
- (d) Show that Legendre polynomials are mutually orthogonal with respect to a scalar product defined as integration over the interval  $[-1, 1]$ , and their norm is  $\sqrt{2/(2l + 1)}$ , i.e.

$$\int_{-1}^1 P_l(\xi) P_{l'}(\xi) d\xi = \frac{2}{2l + 1} \delta_{ll'}. \quad (3)$$

- (e) Now we return to the general case of Legendre's differential equation. Show that for  $m \leq l$  the functions

$$P_l^m(\xi) = (1 - \xi^2)^{\frac{m}{2}} \frac{d^m}{d\xi^m} P_l(\xi) \quad (4)$$

are solutions to (1). They are called *associated Legendre functions of the first kind*.

### [2] The Orbital Angular Momentum Eigenvalue Problem (20 points)

In a previous homework you have shown that the angular momentum operators  $L_z$  and  $L^2$  commute, and you have found explicit *coordinate space* representations for them. Find the common eigenfunctions, i.e. solve the two eigenvalue equations<sup>1</sup>

$$L_z Y(\theta, \phi) = m \hbar Y(\theta, \phi), \quad (5)$$

$$L^2 Y(\theta, \phi) = \lambda \hbar^2 Y(\theta, \phi) \quad (6)$$

---

<sup>1</sup>It is customary to write powers of  $\hbar$  explicitly in this eigenvalue problem.

by choosing a separation ansatz  $Y(\theta, \phi) = \Phi(\phi)\Theta(\theta)$ . The functions  $Y(\theta, \phi)$  in proper normalization (discussed later) are called *spherical harmonics*.

*Hint: First solve for  $\Phi(\phi)$  (what are the allowed values for  $m$ ?) and then show that the equation for  $\Theta$  reduces to Legendre's differential equation from problem [1].*

**[3] Ground State Splitting for the Double Harmonic Oscillator (20 points)**

Consider a particle of mass  $m$  in a double harmonic oscillator potential  $V(x) = \frac{1}{2}m\omega^2(|x| - a)^2$  where  $a$  is the parameter determining the separation of the two harmonic oscillator minima.

(a) We choose trial functions

$$\psi_{\pm}^n = N_{\pm}^n [\psi_n(x - a) \pm \psi_n(x + a)] \quad (7)$$

as discussed in section III.4, where the  $\psi_n$  are the usual harmonic oscillator eigenfunctions. Calculate the values of the functional  $\langle H \rangle[\psi_{\pm}^0]$  for the case  $n = 0$ . As you know they are approximations to the energies of the true ground state and first excited state. *Hint: You can leave integrals for which you do not find simple expressions.*

(b) In the asymptotic limit  $a \rightarrow \infty$  the integrals you obtained in (a) should evaluate to simple expressions. Show that the leading terms in this limit are

$$\langle H \rangle[\psi_{\pm}^0] = \frac{1}{2}\hbar\omega \mp \frac{\alpha}{\sqrt{\pi}}e^{-\alpha^2} \quad (8)$$

in terms of the dimensionless parameter  $\alpha = \sqrt{m\omega/\hbar}x$ . Thus which trial function,  $\psi_+^0$  or  $\psi_-^0$ , is the approximation for the ground state?

**[4] Change of Basis in Hilbert Space (20 points)**

Consider a Hilbert space  $\mathcal{H}$  and let  $|\phi_i\rangle$  and  $|\psi_i\rangle$  be two sets of orthonormal basis vectors with index  $i$  running in some countable set  $I$ .<sup>2</sup> Let  $S$  be a matrix that transforms one basis into the other one by

$$|\phi_i\rangle = \sum_j S_{ij}|\psi_j\rangle \quad (9)$$

for  $i \in I$ .

- (a) Express the coefficients  $S_{ij}$  of  $S$  solely through the states  $|\phi_i\rangle$  and  $|\psi_i\rangle$ ,  $i \in I$ .
- (b) Show that  $S$  is a unitary matrix.
- (c) Let  $A$  be a linear operator on  $\mathcal{H}$ . Given  $S$  and the matrix elements  $A_{ij} = \langle \psi_i | A | \psi_j \rangle$  with respect to the basis  $|\psi_i\rangle$ , calculate the matrix elements  $\tilde{A}_{ij}$  of  $A$  with respect to the basis  $|\phi_i\rangle$ .

---

<sup>2</sup>Otherwise sums have to be replaced by integrals in the usual way but the results still hold.