

---

---

# Physics 606 — Spring 2017

## Homework 5

Instructor: Rainer J. Fries

Turn in your work by March 9

---

---

### [1] Free Particles as a Limit of a Large Potential Well (35 points)

Consider an infinitely deep potential well of size  $L$  with  $V(\vec{r}) = 0$  for  $-L/2 \leq r_i \leq L/2$  for  $i = 1, 2, 3$ ,  $V(\vec{r}) \rightarrow \infty$  elsewhere. Unlike in II.6.4 we now consider solutions of the Schrödinger equation for a particle of mass  $m$  in the potential  $V(\vec{r})$  with periodic boundary conditions (i.e. for opposite boundary points the value of  $\psi$  and all of its derivatives coincide).

*The potential well with periodic boundary conditions and size  $L \rightarrow \infty$  is a useful approximation of free particles.*

- Find the wave functions (with proper normalizations) that are simultaneous eigenfunctions for the three components of the momentum operator,  $p_x$ ,  $p_y$ ,  $p_z$ , together with their eigenvalues. Demonstrate that they are also energy eigenstates of the Hamilton operator and give their energy eigenvalues. Show that for  $L \rightarrow \infty$  the eigenvalues and eigenstates of free particles (albeit with different normalizations) are recovered.
- Introduce a quantum phase space density  $\rho$  by counting the number of eigenstates in a phase space volume  $V_p = L^3 \Delta p_x \Delta p_y \Delta p_z$  and dividing by  $V_p$ . What is the value of  $\rho$ ? Thus what is the average phase space volume occupied by an individual eigenstate?  
*This is an important result for statistical quantum mechanics.*
- Introduce a density  $\sigma = \Delta N / \Delta E$  of eigenstates in the energy spectrum by counting the number of states  $\Delta N$  in an energy interval  $\Delta E$ . Calculate  $\sigma$  as a function of energy  $E$  for large  $E$ .

### [2] Translationally Invariant Systems (30 points)

- Consider the unitary operator

$$\mathbf{U}_a = e^{-\frac{i}{\hbar} \mathbf{p} a} \quad (1)$$

for translations by  $a$  in one dimension where  $\mathbf{p}$  is the momentum operator. Show that its eigenvalues cover the complete unit circle in the space  $\mathbb{C}$  of complex numbers, and that they can be parameterized by  $e^{-\frac{i}{\hbar} K a}$  where the  $K$  are *eigenvalues to the momentum operator*  $\mathbf{p}$ , restricted to  $-\frac{\hbar}{2a} \leq K \leq \frac{\hbar}{2a}$ . What are the corresponding eigenfunctions? What is the degeneracy of each eigenvalue? Is it countable?

*This range for  $K$  is called the first Brillouin zone.*

- Show that the space of eigenfunctions of  $\mathbf{U}_a$  for a fixed eigenvalue (given by the momentum eigenvalue  $K$  as above) can be written in the form

$$\psi_K(x) = e^{\frac{i}{\hbar} K x} u(x) \quad (2)$$

where  $u$  is a square-integrable, periodic function with period  $a$ , i.e.  $u(x + a) = u(x)$ . *Eigenfunctions of the form (2) are called Bloch functions. They play an important role in crystals and other periodic systems.*

[3] **Galilei Boosts** (35 points)

- (a) Recall that a Galilei boost with velocity  $\vec{w}$  acts on a wave function as

$$\psi(\vec{r}, t) \mapsto e^{\frac{i}{\hbar}(m\vec{w}\cdot\vec{r}-\frac{1}{2}mw^2t)}\psi(\vec{r}-\vec{w}t, t). \quad (3)$$

Show that boosts in  $x$ -,  $y$ - and  $z$ -direction can be represented by unitary operators

$$D_{w_i} = e^{\frac{i}{\hbar}K_i w_i} \quad (4)$$

$i = 1, 2, 3$ , with Hermitian generators

$$K_i = mr_i - p_i t. \quad (5)$$

Here  $\vec{r}$  and  $\vec{p}$  are the position and momentum operators for a particle of mass  $m$  and  $t$  is time.

*Hint: Baker-Campbell-Hausdorff*

- (b) For a system with potential energy  $V = 0$  compute the commutators of the boost generators  $K_i$  with the other generators of the Galilei group discussed so far:

$$[K_i, K_j], \quad [K_i, p_j], \quad [K_i, H] \quad (6)$$

for  $i, j = 1, 2, 3$ .

*The set of generators with the commutators as a Lie product is called the Galilei algebra.*

- (c) Let  $D_{\vec{w}_1}, D_{\vec{w}_2}$  be the unitary operators representing boosts by velocities  $\vec{w}_1, \vec{w}_2$ , respectively, and let  $D_{\vec{a}}$  represent a spatial translation by  $\vec{a}$ . Show that  $D_{\vec{w}_2}D_{\vec{w}_1} = D_{\vec{w}_2+\vec{w}_1}$ , i.e. the operators from (a) establish a true (non-projective) representation of boosts alone as a subgroup of  $\mathcal{G}_+^+$ . Now consider a spatial translation followed by a boost, once as a product of the individual operators  $D_{\vec{w}_1}D_{\vec{a}}$ , and once as the single operator  $D_{\vec{w}_1\oplus\vec{a}} = e^{\frac{i}{\hbar}(K\cdot\vec{w}_1-\vec{p}\cdot\vec{a})}$  that represents it. From a comparison of the two conclude whether the representation of the Galilei group discussed here is projective.