

Constraining the in-medium nucleon-nucleon cross section from the width of nuclear giant dipole resonance

Rui Wang,^{1,2} Zhen Zhang,³ Lie-Wen Chen,⁴ Che Ming Ko, and Yu-Gang Ma^{1,2}

¹Key Laboratory of Nuclear Physics and Ion-beam Application (MOE), Institute of Modern Physics, Fudan University, Shanghai 200433, China

²Shanghai Institute of Applied Physics, Chinese Academy of Sciences, Shanghai 201800, China

³Sino-French Institute of Nuclear Engineering and Technology, Sun Yat-Sen University, Zhuhai 519082, China

⁴School of Physics and Astronomy and Shanghai Key Laboratory for Particle Physics and Cosmology, Shanghai Jiao Tong University, Shanghai 200240, China

We have used the lattice Hamiltonian (LH) method to solve the Boltzmann-Uheling-Ulenbeck (BUU) transport equation with the binary collisions in the collision term treated via the stochastic approach [1]. By incorporating the stochastic NN collision term into the previous lattice-Hamiltonian Vlasov (LHV) framework [2], we have solved the BUU equation with nuclear mean field obtained from the N3LO Skyrme pseudopotential through the LH method. With the use of a sufficiently large number of




Fig. 1. The GDR width of ^{208}Pb from LBUU calculations for different values of α in $\sigma_{NN}^* = \sigma_{NN}^{\text{free}} \exp\left[-\alpha \frac{\rho/\rho_0}{1+(T_{\text{c.m.}}/T_0)^2}\right]$. The horizontal line represents the RCNP experimental value of 4.0 MeV [3]. The inset shows the strength function with $\alpha=1.8$ (solid line) and the shifted one (dash-dotted line) to match the experimental GDR peak energy.

test particles, the present lattice-Hamiltonian BUU (LBUU) method treats the Pauli blocking in the collision term of BUU equation with very high precision and thus significantly increases the accuracy in solving the BUU equation. From the accurately calculated giant dipole resonance (GDR) width of ^{208}Pb , we have found that it depends strongly on the magnitude of the in-medium NN cross section σ_{NN}^* , and the experimentally measured GDR width of ^{208}Pb from the $^{208}\text{Pb}(\vec{p}, \vec{p}')$ reaction at RCNP [3] can only be reproduced with a NN cross section that is significantly reduced in nuclear medium as shown in Fig.1. The large medium reduction of the free-space NN scattering cross section $\sigma_{NN}^{\text{free}}$ raises challenges to microscopic calculations based on realistic NN interactions. Also, the effects of such a large medium reduction of $\sigma_{NN}^{\text{free}}$ on the widths of other modes of giant resonances in nuclei and on the dynamics of HICs need to be studied as it may significantly affect the extracted information on the properties of nuclear matter at various densities.

[1] R. Wang, Z. Zhang, L.W. Chen, C.M. Ko, and Y.G. Ma, Phys. Lett. B **807**, 135532 (2020).

[2] R. Wang, L.-W. Chen, Z. Zhang, Phys. Rev. C **99**, 044609 (2019).

[3] A. Tamii *et. al.*, Phys. Rev. Lett. **107**, 062502 (2011).